AMENDMENTS TO THE CLAIMS

Claims 1-12: (canceled)

13. (currently amended) An adduct comprising MgCl₂, ethanol and a Lewis base (LB) different from water, said adduct further comprising a fusion enthalpy lower than 100 J/g, and formula MgCl₂•(EtOH)_n(LB)_p, wherein n is from 2 to 6 and p is 0<p/(n+p)≤0.1, where the Lewis base is selected from ethers, esters, compounds of formula RX_m, and combinations thereof, wherein RX_m is selected from the group consisting of methanol, propanol, isopropanol, n-butanol, sec-butanol, tert-butanol, pentanol, 2-methyl-1-pentanol, 2-ethyl-1-hexanol, phenol, 4-methyl-1-phenol, 2,6-dimethyl-1-phenol, cyclohexanol, cyclopentanol, ethylene glycol, propylene glycol, 1,4-butanediol, glycerine, mannitol, polyvinyl-alcohol, acetonitrile, ethylenediamine, 3-picoline, triethanolamine, triethylamine, and diisopropylamine.

(previously presented) The adduct according to claim 13, wherein p is 0
p/(n+p)≤0.0125.

15. (canceled)

16. (canceled)

17. (canceled)

 (previously presented) A catalyst component for polymerizing at least one olefin comprising a product of a reaction between a transition metal compound and the adduct according to claim 13.

19. (previously presented) The catalyst component according to claim 18, wherein the transition metal compound is selected from at least one titanium compound comprising formula Ti(OR)_nX_{y-an} wherein n is between 0 and y; y is a valence of titanium; X is halogen; and R is an alkyl radical comprising 1-8 carbon atoms, or COR, wherein R is a hydrocarbon group comprising from 1 to 20 carbon atoms.

 (previously presented) The catalyst component according to claim 19, wherein the titanium compound is selected from TiCl₃, TiCl₄, Ti(OBu)₄, Ti(OBu)₂Cl₃, Ti(OBu)₂Cl₃, and Ti(OBu)₃Cl.

21. (previously presented) The catalyst component according to claim 18, wherein the reaction between the transition metal compound and the adduct is carried out in presence of an electron donor compound.

22. (previously presented) The catalyst component according to claim 21, wherein the electron donor is selected from esters, ethers, amines, and ketones.

 (previously presented) A catalyst for polymerizing at least one olefin comprising a product of a reaction between the catalyst component according to claim 19, and an aluminum alkyl compound.

24. (previously presented) A process for polymerizing at least one olefin of formula CH₂=CHR, wherein R is hydrogen or a hydrocarbon radical comprising 1-12 carbon atoms, carried out in presence of the catalyst according to claim 23.

25. (canceled)

26. (canceled)

27. (currently amended) The adduct of claim 4613, wherein RX_m is selected from the group consisting of phenol, 4-methyl-1-phenol, 2,6-dimethyl-1-phenol, ethylene glycol, propylene glycol, 1,4-butanediol, glycerine, mannitol, polyvinyl-alcohol, acetonitrile, ethylenediamine, 3-picoline, triethanolamine, triethylamine, and diisopropylamine.